

# Can a dust dominated universe have accelerated expansion?

Håvard Alnes<sup>1</sup>, Morad Amarzguioui<sup>2</sup> and Øyvind Grøn<sup>1,3</sup>

<sup>1</sup> Department of Physics, University of Oslo, PO Box 1048 Blindern, 0316 Oslo, Norway

<sup>2</sup> Institute of Theoretical Astrophysics, University of Oslo, PO Box 1029 Blindern, 0315 Oslo, Norway

<sup>3</sup> Oslo College, Faculty of Engineering, Cort Adelersgt. 30, 0254 Oslo, Norway

**Abstract.** Recently, there has been suggestions that the apparent accelerated expansion of the universe is due not to a cosmological constant, but rather to inhomogeneities in the distribution of matter. In this work, we investigate a specific class of inhomogeneous models that can be solved analytically, namely the dust-dominated Lemaitre-Tolman-Bondi universe models. We show that they do not permit accelerated cosmic expansion.

**Keywords:** inhomogeneous universe models – dark energy – accelerated expansion

## 1. Introduction

Kolb and coworkers [1] argued recently that growing perturbations on a scale larger than the Hubble length may lead to accelerated expansion as observed from the center of the perturbation. This was then followed up by Wiltshire [2] and Carter et al. [3], who considered a dust dominated universe model where the observed universe is an underdense bubble in an Einstein-de Sitter universe. They calculated the luminosity-redshift relationship for their model and found very small deviations from the standard  $\Lambda$ CDM model. Thus, they were able to obtain a good fit to the Hubble diagram of type Ia supernovae.

However, critical papers have appeared. We will now proceed to give a brief overview over some of these papers and the arguments therein. By considering a universe model where sub-Hubble perturbations are absent, Flanagan [4] argues that the contributions from the super-Hubble perturbations to the value of the deceleration parameter are so small that they cannot be responsible for the acceleration of the universe. Geshnizjani et al. [5] argue that to second order in spatial gradients the super-Hubble perturbations only amount to a renormalization of local spatial curvature, and thus cannot account for the negative deceleration parameter. Making an expansion to Newtonian order in potential and velocity, but taking into account fully non-linear density inhomogeneities, Siegel and Fry [6] obtain similar results, and conclude that inhomogeneity contributions cannot mimic the effects of dark energy or induce an accelerated expansion.

One of the strongest and most convincing critiques of the conclusions of Kolb et al. is that of Hirata and Seljak [7]. They argue from the Raychaudhuri equation that in a dust dominated universe there must be a non-vanishing vorticity in order to obtain a negative deceleration parameter. Then they showed that the perturbations considered in [1] have vanishing vorticity and hence cannot lead to accelerated expansion. In yet another critique of the work of Kolb et al., Räsänen [8] also claims that their conclusions are ruled out. The basis for this conclusion was an analysis of the model proposed in [1] by applying the Buchert formalism for backreaction [9]. This conclusion is also supported in works by Giovannini [10, 11], where the author performs a general analysis of the class of models considered in [1]. He finds that dust-dominated models cannot have accelerated expansion, and hence, the conclusion of Kolb et al. cannot be correct.

Several of the arguments against accelerated expansion induced by inhomogeneities in matter are based upon perturbation calculations. However, as noted in [5], one way of evading these arguments is through non-perturbative effects. This is what we will investigate in the present work. We want to know if Einstein's field equations permit inhomogeneities to change the positive deceleration parameter of a dust-dominated homogeneous universe model to a negative value.

Unfortunately, a general analysis incorporating all possible inhomogeneous models is not feasible, so we will instead concentrate on one specific class of such models, namely the dust-dominated, spherically symmetric models. Moffat suggested in [12]

that it might be possible to find solutions with accelerated expansion. Considering the field equations for these models, an expression for the local deceleration parameter was derived, and it was claimed that this might indeed allow for negative values. Later, in [13], this claim was moderated considerably. Here it was claimed that the *volume averaged* deceleration parameter, and not the local deceleration parameter, could be negative. No explicit solution of the field equation were actually found in these analyses. Instead, the conclusions were reached by investigating the field equations directly.

In this work, we will consider the same class of spherically symmetric, inhomogeneous universe models and present a general solution to the field equations. We then show that no such model can have an accelerated expansion in the sense of a negative local deceleration parameter.

The structure of this paper is as follows. In Sect. 2 we present the field equations for a general spherically symmetric, inhomogeneous universe model. In Sect. 3 we discuss the properties of these models and whether they can have accelerated expansion. Finally, in Sect. 4 we give a brief summary of our work and present our conclusion.

## 2. Spherically symmetric, inhomogeneous universe models

The spherically symmetric, inhomogeneous universe models are described by the Lemaitre-Tolman-Bondi (LTB) space-time [14, 15, 16]. The line element can be written as

$$ds^2 = -dt^2 + X^2(r, t)dr^2 + R^2(r, t)d\Omega^2, \quad (1)$$

where  $X(r, t)$  and  $R(r, t)$  are general function to be determined by the field equations and boundary and initial conditions. They can be thought of as generalized, position dependent scale factors in the radial and transverse directions. The Einstein field equations are

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \kappa T_{\mu\nu}, \quad (2)$$

where  $\kappa = 8\pi G$ .

We assume the universe to be occupied by an ideal fluid with energy density  $\rho(r, t)$  and an isotropic pressure  $p(r, t)$ . In the comoving coordinates defined in Eq. (1), the energy-momentum tensor of the fluid can be written as  $T_{\mu\nu} = \text{diag}(\rho, p, p, p)$ . A relation between  $X$  and  $R$  can be found by solving the  $t$ - $r$  component of the field equations, for which the right-hand side vanishes. The relation one finds is

$$X(r, t) = \frac{R'(r, t)}{f(r)}, \quad (3)$$

where  $f(r)$  is an arbitrary function of  $r$  only. Throughout this paper, we will use a  $' = d/dr$  to denote differentiation with respect to  $r$  and  $\dot{\phantom{x}} = d/dt$  for differentiation with respect to  $t$ .

Following Moffat [12, 13], we define two "Hubble parameters"

$$H_{\perp} = \frac{\dot{R}}{R} \quad \text{and} \quad H_r = \frac{\dot{R}'}{R'}, \quad (4)$$

which are a measure of the local expansion rate in the perpendicular and radial directions respectively, and the deceleration parameter

$$q_{\perp} = -\frac{1}{H_{\perp}^2} \frac{\ddot{R}}{R}. \quad (5)$$

Using these definitions, the time-time and space-space components of the field equations take the form

$$H_{\perp}^2 + 2H_r H_{\perp} - \frac{\beta}{R^2} - \frac{\beta'}{RR'} = \kappa\rho \quad (6)$$

and

$$-6H_{\perp}^2 q_{\perp} + 2H_{\perp}^2 - 2\frac{\beta}{R^2} - 2H_r H_{\perp} + \frac{\beta'}{RR'} = -\kappa(\rho + 3p). \quad (7)$$

where  $\beta \equiv f^2 - 1$ . These are the generalization of the Friedmann equations in a homogeneous universe to a spherically symmetric, inhomogeneous universe. Note that the function  $\beta(r)$  is often written as  $\beta(r) = 2E(r)r^2$ , where  $E(r)$  determines the local curvature radius.

### 3. Expansion rate of dust-dominated LTB models

From now on we will assume the cosmic fluid to be pressure-less matter. Adding Eqs. (6) and (7), we then obtain the following expression for the deceleration parameter

$$q_{\perp} = \frac{1}{2} - \frac{\beta}{2\dot{R}^2}. \quad (8)$$

The condition for accelerated expansion now takes the form

$$\beta > \dot{R}^2 > 0 \quad \text{or} \quad f^2 > 1 + \dot{R}^2. \quad (9)$$

Substituting the definition for the deceleration parameter in Eq. (5) into Eq. (8), we arrive at the expression

$$2R\ddot{R} + \dot{R}^2 = \beta. \quad (10)$$

A single integration of this equation yields

$$R\dot{R}^2 = \beta R + \alpha \quad \text{or} \quad H_{\perp}^2 = \frac{\beta}{R^2} + \frac{\alpha}{R^3}, \quad (11)$$

where the function  $\alpha(r)$  enters as an integration “constant” and depends only on the radial coordinate  $r$ . Comparing this equation with the ordinary Friedmann equation for homogeneous models, we see that the dynamical effects of  $\beta$  and  $\alpha$  are similar to those of curvature and dust, respectively. Therefore,  $\alpha(r)$  is regarded as the gravitational mass function, and one often chooses  $\alpha(r) \propto r^3$ . Note, however, that  $\alpha$  does not appear in the energy-momentum tensor.

From Eqs. (9) and (11), we find that  $\alpha$  and  $\beta$  must satisfy the inequality

$$-\beta R < \alpha(r) < 0 \quad (12)$$

in order for the deceleration parameter to be negative. Thus, it appears that the LTB models seem to allow for accelerated expansion even for dust-dominated universe models,

as long as  $\alpha$  and  $\beta$  are chosen appropriately. The dynamical effect of  $\alpha$  would then correspond to that of dust with negative energy density in a homogeneous universe model. It should be noted that the inequality above forbids accelerated expansion in a “big bang” model where the scale factor has the initial value  $R(0, r) = 0$ , which implies that  $\alpha(r) > 0$ . However, this initial condition may not be physically realistic. The universe may have started with a finite scale factor, or maybe has collapsed and reached a finite minimum radius before expanding again. In such models accelerated expansion does not seem to be forbidden. However, as we will show shortly, such solutions are unphysical. First, we find a general solution of the field equation and show how such an accelerated solution behaves.

We introduce a conformal time  $\eta$  defined via the differential relation  $\beta^{1/2}dt = R d\eta$ . This allow us to integrate Eq. (11) to give a parametric solution in terms of  $\eta$ . The solutions can be grouped into three different classes according to the sign of the local curvature  $\beta$ . Choosing  $t_0 = \eta_0 = 0$ , the three classes are

$$\begin{aligned} R &= \frac{\alpha}{2\beta}(\cosh \eta - 1) + R_0 \left[ \cosh \eta + \sqrt{\frac{\alpha + \beta R_0}{\beta R_0}} \sinh \eta \right] \\ \sqrt{\beta}t &= \frac{\alpha}{2\beta}(\sinh \eta - \eta) + R_0 \left[ \sinh \eta + \sqrt{\frac{\alpha + \beta R_0}{\beta R_0}} (\cosh \eta - 1) \right] \quad , \beta > 0 \end{aligned} \quad (13)$$

for positive  $\beta$ ,

$$\begin{aligned} R &= \frac{\alpha}{2|\beta|}(1 - \cos \eta) + R_0 \left[ \cos \eta + \sqrt{\frac{\alpha + \beta R_0}{|\beta| R_0}} \sin \eta \right] \\ \sqrt{|\beta|}t &= \frac{\alpha}{2|\beta|}(\eta - \sin \eta) + R_0 \left[ \sin \eta + \sqrt{\frac{\alpha + \beta R_0}{|\beta| R_0}} (1 - \cos \eta) \right] \quad , \beta < 0 \end{aligned} \quad (14)$$

for negative  $\beta$ , and finally


$$R = \left( R_0^{3/2} + \frac{3}{2}\sqrt{\alpha}t \right)^{2/3} \quad (15)$$

for vanishing  $\beta$ . We allow the initial size of the universe to be non-zero, hence the non-vanishing value for  $R_0 \equiv R(t = 0, r)$ . Eqs. (13)-(15) with  $R_0 = 0$  represent the usual form of the LTB solution of the field equations. The scale factor  $R$  as a function of time is shown for some typical parameters in Fig. 1. The top curve represents a solution with accelerating expansion.

In order for a dust-dominated solution in the LTB space-time to undergo an epoch of accelerated expansion, the inequality in Eq. (12) has to be satisfied. Can a physically realistic solution satisfy this? Differentiating Eq. (11) and comparing the result with Eq. (6) we obtain a relation between the density distribution and  $\alpha$  for this class of models,

$$\kappa\rho = \frac{\alpha'}{R^2 R'} = 3 \frac{\alpha'}{V'} \quad (16)$$

where  $V = R^3$  is a comoving volume. A physically realistic model must have positive density everywhere, and hence, it must have  $\alpha' > 0$ . Furthermore, the angular part of



**Figure 1.** The scale factor for some typical values of  $\alpha$  and  $\beta$

the line element is  $R^2 d\Omega^2$  where  $d\Omega$  is a solid angle element. At the origin,  $r = 0$ , this part of the line element must vanish, and thus  $R(0, t) = 0$ . From Eq. (11) we then get that  $\alpha(0) = 0$ . Since  $\alpha'(r) > 0$ , it follows that  $\alpha(r) \geq 0$  for all  $r$ . But, as we showed above, accelerated expansion is only possible for models with  $\alpha(r) < 0$ . Hence, the dust dominated LTB universe models must have decelerated expansion.

We shall now arrive at the same conclusion by an alternative approach related to the Raychaudhuri equation, which describes the flow of a cosmological fluid in space-time. It can be written as

$$\nabla_\mu a^\mu = \dot{\theta} + \frac{\theta^2}{3} + 2(\sigma^2 - \omega^2) + R_{\mu\nu} u^\mu u^\nu, \quad (17)$$

where  $a^\mu \equiv u^\mu_{;\alpha} u^\alpha$  is the four-acceleration of the fluid,  $\theta \equiv u^\mu_{;\mu}$  is the expansion rate of the fluid and  $\sigma$  and  $\omega$  are the shear and vorticity scalars, respectively. The latter two quantities are defined as

$$\sigma^2 = \frac{1}{2} \sigma_{\mu\nu} \sigma^{\mu\nu} \quad \text{and} \quad \omega^2 = \frac{1}{2} \omega_{\mu\nu} \omega^{\mu\nu} \quad (18)$$

where

$$\sigma_{\mu\nu} = u_{(\mu;\nu)} + a_{(\mu} u_{\nu)} - \frac{1}{3} \theta (g_{\mu\nu} + u_\mu u_\nu) \quad (19)$$

and

$$\omega_{\mu\nu} = u_{[\mu;\nu]} + a_{[\mu} u_{\nu]}. \quad (20)$$

In a comoving reference frame the four-velocity of the fluid can be written as  $u^\mu = [1, 0, 0, 0]$ . The divergence of the four-acceleration will then vanish, and the Raychaudhuri equation can be simplified to

$$0 = \dot{\theta} + \frac{\theta^2}{3} + 2(\sigma^2 - \omega^2) + R_{00}. \quad (21)$$

Analogously to the Hubble parameter in FRW models, we can define an effective local Hubble parameter from the expansion rate of the fluid:

$$H = \frac{1}{3} u^\mu_{;\mu}. \quad (22)$$

This allow us to define an effective deceleration parameter in terms of  $H$  and  $\dot{H}$ :

$$q \equiv -\frac{\dot{H} + H^2}{H^2} = -1 - 3\dot{\theta}/\theta^2. \quad (23)$$

Using the relation  $R_{00} = \frac{\kappa}{2}(\rho + 3p)$ , Eq. (21) can be written

$$\theta^2 q = 6(\sigma^2 - \omega^2) + \frac{3}{2}\kappa(\rho + 3p). \quad (24)$$

While Eq. (24) is valid for any metric, let us now specialize to the case of an LTB model. In [12] it was claimed that the vorticity scalar is non-vanishing for the models considered in the paper, which may permit negative values of  $q$ . We find, however, that the vorticity tensor (20) vanishes for a comoving fluid in the metric (1). This incorporates also the models considered in [12]. For this metric the shear can be written as

$$\sigma^2 = \frac{1}{3}(H_r - H_\perp)^2. \quad (25)$$

Finally, assuming the fluid to be pressure-less, Eq. (24) reduces to

$$\theta^2 q = 2(H_r - H_\perp)^2 + \frac{3}{2}\kappa\rho. \quad (26)$$

This equation shows that the dust dominated LTB-models have  $q \geq 0$ , meaning that they cannot have accelerated expansion.

Although we have now shown that the LTB models cannot have accelerated expansion, this doesn't necessarily mean that they cannot explain the data used to infer this expansion. The first data used to conclude that the expansion seems to be accelerating were the measurements of the luminosities of supernovae of type Ia (SNIa) [17, 18]. Since then newer data have appeared that strengthen this claim even further [19, 20, 21]. However, accelerated expansion follows logically from these data only if one assumes that the universe is homogeneous. The added freedom of having a position dependent expansion in LTB models allows one to explain the data without the need for the expansion to accelerate locally. The explanation would then be that the expansion rate is highest at  $r = 0$  and decreases with distance from the center, since the oldest supernovae are also farthest away. In this case a decelerating universe would seem to be accelerating. C  lerier shows in [22] how such inhomogeneities can mimic the effects of dark energy in supernova observations, when interpreted within the framework of FRW models. More specifically, Iguchi et al. [23] show that models with  $\alpha(r) > 0$  can

reproduce the luminosity distance-redshift relationship of a  $\Lambda$ CDM model up to  $z \sim 1$ , but not for higher redshifts. However, when compared to actual SNIa observations, it is easy to find models that fit the data even better than the best-fit  $\Lambda$ CDM model [24]. In fact, as we show in [24], these models might also be compatible with the CMB power spectrum, even though they do not contain any form of dark energy.

Finally, we wish to stress the point that in this work we have shown that the LTB models cannot have an expansion that is *locally* accelerated. This does not exclude the possibility that there can be a so-called *volume averaged acceleration*, where a scale factor defined via the physical volume of a comoving region is found to have a positive double time derivative. Recent papers discussing the possibilities of this effect explaining the apparent accelerated expansion are Refs. [25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36]. At the present time there does not appear to exist a consensus among cosmologists on this issue.

#### 4. Conclusion

Recently it was claimed by Kolb et al. [1] that inhomogeneous perturbations of FRW universe models can result in accelerated cosmic expansion, and thereby eliminate the need to postulate the existence of the mysterious dark energy. However, several papers have appeared since criticizing this work heavily. These attribute the effects claimed by Kolb et al. to incorrect handling of second order terms in the perturbative expansion. When dealt with correctly, one would then assume that these terms do not lead to accelerated expansion. However, this leaves open the possibility that the full, non-perturbative solutions of the Einstein equations for inhomogeneous models might exhibit accelerated expansion.

In this work we investigated a special class of inhomogeneous models which can be solved analytically, the so-called Lemaitre-Tolman-Bondi (LTB) models. These are the spherically symmetric, inhomogeneous models. We found a general solution to a dust-dominated LTB model, and showed that there exist solutions that appear to have accelerated expansion. However, these solutions turned out to be unphysical. Finally, it was shown that any physically realistic solution of Einstein's equations must have decelerated expansion if it contains only matter.

#### Acknowledgments

MA acknowledges support from the Norwegian Research Council through the project 159637/V30 – “Shedding Light on Dark Energy”.

#### References

- [1] Kolb E W, Matarrese S, Notari A and Riotto A 2005 (*Preprint hep-th/0503117*)
- [2] Wiltshire D L 2005 (*Preprint gr-qc/0503099*)



- [3] Carter B M N, Leith B M, Ng S C C, Nielsen A B and Wiltshire D L 2005 (*Preprint astro-ph/0504192*)
- [4] Flanagan E E 2005 *Phys. Rev.* **D71** 103521 (*Preprint hep-th/0503202*)
- [5] Geshnizjani G, Chung D J H and Afshordi N 2005 *Phys. Rev.* **D72** 023517 (*Preprint astro-ph/0503553*)
- [6] Siegel E R and Fry J N 2005 *Astrophys. J.* **628** L1–L4 (*Preprint astro-ph/0504421*)
- [7] Hirata C M and Seljak U 2005 *Phys. Rev.* **D72** 083501 (*Preprint astro-ph/0503582*)
- [8] Rasanen S 2006 *Class. Quant. Grav.* **23** 1823–1835 (*Preprint astro-ph/0504005*)
- [9] Buchert T 2000 *Gen. Rel. Grav.* **32** 105–125 (*Preprint gr-qc/9906015*)
- [10] Giovannini M 2006 *Phys. Lett.* **B634** 1–4 (*Preprint hep-th/0505222*)
- [11] Giovannini M 2005 *JCAP* **0509** 009 (*Preprint astro-ph/0506715*)
- [12] Moffat J W 2005 (*Preprint astro-ph/0504004*)
- [13] Moffat J W 2006 *JCAP* **0605** 001 (*Preprint astro-ph/0505326*)
- [14] Lemaitre G 1933 *Ann. Soc. Sci. Brux.* **A53** 51–85
- [15] Tolman R C 1934 *Proc. Nat. Acad. Sci.* **20** 169–176
- [16] Bondi H 1947 *Mon. Not. Roy. Astron. Soc.* **107** 410–425
- [17] Riess A G *et al.* 1998 *Astron. J.* **116** 1009 (*Preprint astro-ph/9805201*)
- [18] Perlmutter S *et al.* 1999 *Astrophys. J.* **517** 565 (*Preprint astro-ph/9812133*)
- [19] Tonry J L *et al.* 2003 *Astrophys. J.* **594** 1 (*Preprint astro-ph/0305008*)
- [20] Knop R A *et al.* 2003 *Astrophys. J.* **598** 102 (*Preprint astro-ph/0309368*)
- [21] Riess A G *et al.* (Supernova Search Team) 2004 *Astrophys. J.* **607** 665–687 (*Preprint astro-ph/0402512*)
- [22] Celerier M N 2000 *Astron. Astrophys.* **353** 63–71 (*Preprint astro-ph/9907206*)
- [23] Iguchi H, Nakamura T and Nakao K 2002 *Prog. Theor. Phys.* **108** 809–818 (*Preprint astro-ph/0112419*)
- [24] Alnes H, Amarzguioui M and Gron O 2006 *Phys. Rev.* **D73** 083519 (*Preprint astro-ph/0512006*)
- [25] Kolb E W, Matarrese S and Riotto A 2005 (*Preprint astro-ph/0506534*)
- [26] Buchert T 2005 *Class. Quant. Grav.* **22** L113–L119 (*Preprint gr-qc/0507028*)
- [27] Buchert T 2006 *Class. Quant. Grav.* **23** 817–844 (*Preprint gr-qc/0509124*)
- [28] Buchert T, Larena J and Alimi J M 2006 (*Preprint gr-qc/0606020*)
- [29] Nambu Y and Tanimoto M 2005 (*Preprint gr-qc/0507057*)
- [30] Ishibashi A and Wald R M 2006 *Class. Quant. Grav.* **23** 235–250 (*Preprint gr-qc/0509108*)
- [31] Mansouri R 2005 (*Preprint astro-ph/0512605*)
- [32] Mansouri R 2006 (*Preprint astro-ph/0601699*)
- [33] Kasai M, Asada H and Futamase T 2006 *Prog. Theor. Phys.* **115** 827–832 (*Preprint astro-ph/0602506*)
- [34] Paranjape A and Singh T P 2006 (*Preprint astro-ph/0605195*)
- [35] Parry M F 2006 (*Preprint astro-ph/0605159*)
- [36] Kai T, Kozaki H, Nakao K, Nambu Y and Yoo C 2006 (*Preprint gr-qc/0605120*)